



Research Article

QUALITY CHARACTERIZATION OF MANDARIN (*Citrus reticulata* Blanco) DURING STORAGE

PIPPAL RAHUL¹, GUPTA RAJESH², KUCHI VENKATA SATISH^{*3} AND RANA SWETAL⁴

^{1,2,4}Department of Postharvest Management, KNK College of Horticulture, RVSKVV, Mandsaur, Madhya Pradesh, India

³Department of Postharvest Technology, Faculty of Horticulture, BCKV, Mohanpur, 741252, West Bengal, India

*Corresponding Author: Email-newmoon_9@yahoo.com

Received: April 27, 2016; Revised: May 10, 2016; Accepted: May 11, 2016; Published: September 06, 2016

Abstract- Quality characterization of mandarin was done with the objective to analyze and find out best treatment among different combinations of pre-treatments and packaging during storage at ambient conditions (19 to 25°C). Twenty treatment combinations consisting of four types of chemicals (2, 4- Dichlorophenoxyacetic acid 50 and 100 ppm, Gibberellic acid 50 and 100 ppm) and three types of packaging materials (Corrugated boxes, perforated polythene and newspaper) and their combinations were used in the study. Maximum values for titrable acidity (0.624%), TSS (13.12° Brix), total sugars (9.56%) and ascorbic acid (14.17 mg/100 ml juice) were obtained with 2,4-D 100 ppm with perforated polythene on 20th day of storage. Maximum scores for sensory quality were observed in the treatment combination with 2, 4-D 100 ppm and corrugated fibre board boxes.

Keywords- Ascorbic acid, Mandarin, TSS, 2,4-D and Overall acceptability.

Citation: Pippal Rahul, et al., (2016) Quality Characterization of Mandarin (*Citrus reticulata* Blanco) During Storage. International Journal of Agriculture Sciences, ISSN: 0975-3710 & E-ISSN: 0975-9107, Volume 8, Issue 33, pp.-1697-1701.

Copyright: Copyright©2016 Pippal Rahul, et al., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Academic Editor / Reviewer: Jayaprakash S. M.

Introduction

Citrus reticulata Blanco are commonly known as mandarins and their rind is loosely attached to the juicy sacs. Mandarin is native to tropical and subtropical region of South-East Asia. They are known to cure fever. They come to harvest in 32-36 weeks after the fruit set. Mandarins are harvested based on the skin colour. When the surface of the fruit turns 25% or more yellow orange colour, they can be harvested. The peel colour improved with advancement of maturity [1]. It is available from mid October to January end in Malwa region of Madhya Pradesh. If we use appropriate packaging materials and suitable chemicals for storing of the fresh fruits, we can increase the shelf-life of mandarin. They can be transported to local and short distant markets with minimum physiological loss in weight. Market value of citrus fruits is controlled by its quality, which is dependent on its external and internal characteristics. Maintaining quality is one of the main factors in increasing sales. Huge harvest of produce during peak harvesting season creates glut and the growers are compelled to sell it in the local or nearby markets at throwaway rates. Hence, mandarin packaging using various packaging materials, storage and value addition is a solution to benefit the farmer of the Malwa region of Madhya Pradesh. Fruit colour retained greenness of buttons in Nagur mandarin with 2,4-D treatment [2]. Therefore, keeping these points in view a study on packaging and storage of mandarin is planned to access the quality characteristics of mandarin when subjected to different treatments.

Materials and methods

The experiment titled, "Quality characterization of mandarin (*Citrus reticulata* Blanco) during storage" was conducted at the Department of Post Harvest Management, College of Horticulture, Mandsaur, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior during 2010-2011. For different treatment combinations, one level of variety, five levels of pre-treatments and four levels of

methods of packaging were used. Thus, in the present investigation a total of 20 treatment combinations on storage were made. The experimental details and various treatment combinations are presented in [Table-1].

Table-1 Experimental details

Variety	Nagpur
Pre-treatment	1. Untreated (Without any chemical treatment) (T ₀)
	2. Dipping in 2,4-D 50 ppm (T ₁)
	3. Dipping in 2,4-D 100 ppm (T ₂)
	4. Dipping in GA ₃ 50 ppm (T ₃)
	5. Dipping in GA ₃ 100 ppm (T ₄)
Packaging materials	1. Untreated (Without any packaging material) (P ₀)
	2. Corrugated fiber box (P ₁)
	3. Perforated polythene (P ₂)
	4. Individual wrapping with newspaper (P ₃)
Total No. of treatment combinations	20
No. of replications	3
Quantity of fruits per pack	2 kg
Total number of treatments	20×3 = 60
Experimental design	Factorial CRD
Observations recorded at	0, 5 th , 10 th , 15 th and 20 th day of storage

Evenly sized, uniform and fully matured fruits of mandarin cv. Nagpur were harvested from the field of a farmer Shri Rajendra Patidar of Jaora, Ratlam (M.P.)

and brought to the laboratory for the purpose of experiment. A sample of 2 kg mandarin fruits/pack was weighed before packaging and storage. For a single replication 40 kg fruits were used in the experiment. Damaged, diseased, and immature fruits were sorted out. Selected fruits were washed with the help of running tap water. Thus for three replications a total of 120 kg mandarin was used for the experiment.

Preparation of chemical solution

For preparation of 50 and 100 ppm 2, 4-D solution, 0.05 and 0.1 g crystals of 2, 4-D were dissolved in 10 ml of ethanol respectively. Thereafter, 1 litre of distilled water was added to it. Similarly, for 50 and 100 ppm GA₃ solution, 0.05 and 0.1 g crystals of GA₃ were dissolved in 10 ml of ethanol respectively. Thereafter, 1 litre of distilled water was added to it.

Pretreatment

The washed fruits were dipped in above prepared chemical solutions for five minutes. After pre-treatment the fruits were dried at room temperature for one hour.

Packaging and Storage

After moisture removal the mandarin fruits were packed in different packaging material (corrugated boxes, perforated polythene and newspaper) @ 2 kg/pack. After packaging, the fruits were stored at ambient room temperature (16-22°C) and relative humidity (70-80%). The treated fruits were subjected to various physico-chemical observations at 0, 5th, 10th, 15th and 20th day of storage as described in succeeding sub-sections.

Organoleptic characteristics of fresh and stored mandarin

The mandarin fruit is stored over a period of 20 days were subjected to organoleptic evaluation by a panel of six judges following hedonic rating tests as described by [3]. The mandarin fruits were evaluated for colour and overall acceptability. The characters with mean scores of zero or more out of 10 marks were considered acceptable.

Biochemical characteristics of fresh and stored mandarin

Acidity

Acidity (citric acid) was determined by diluting the known weight of fresh juice of mandarin fruit sample (sample was weight and ground well with a pestle and mortar) and titrating the sample against standard 0.1N NaOH using phenolphthalein as the indicator. Appearance of light pink colour denotes the end point. Acidity had been calculated by using following formula and expressed in percent.

$$\text{Acidity (\%)} = \frac{1 \times \text{Normality of NaOH} \times \text{Equivalent weight of acid} \times \text{Titer}}{10 \times \text{Weight of the sample}} \times 100$$

Ascorbic acid

Ascorbic acid content of fresh and stored fruits was determined by 2, 6-dichlorophenol indophenol dye solution method [4] and expressed in mg ascorbic acid (vitamin C) per 100 ml of fruit juice. Formula for calculating ascorbic acid content was given under

$$\text{Ascorbic acid (mg/100 ml)} = \frac{\text{Titer} \times \text{Dye equivalent} \times \text{Dilution}}{\text{Weight of the sample}} \times 100$$

Total sugars

Total sugar content was estimated by measuring the absorbance at 620 nm as suggested by [3] and expressed in percentage basis.

Total soluble solids (TSS)

TSS content was measured at room temperature with the help of Abbe refractometer (Fuzhou, made in China).

Temperature and relative humidity

Ambient temperature was determined with the help of minimum and maximum thermometer (Zeal -Made in England, Capacity - 0°C to 50°C). Relative humidity was measured through digital hygrometer (make Vista Biocell Pvt. Ltd. New Delhi, INDIA). Dry bulb and wet bulb temperatures were also used to calculate the relative humidity with the help of psychometric chart.

Statistical Analysis

Completely Randomized Design was employed to test the significance of variation in the data obtained as suggested by [5].

Results and Discussion

Chemical characteristics

Quality characteristics of fresh mandarin viz., acidity, ascorbic acid, reducing sugars, total sugars and total soluble solids were determined shown in [Table-2].

Table-2 Quality characteristics of fresh mandarin

S. No.	Characteristics	Value
1.	Acidity (%)	0.89
2.	Ascorbic acid (mg/100ml)	17.4
3.	Total sugars (%)	10.3
4.	T.S.S. (°Brix)	7.4
5.	Colour	9.5
6.	Overall acceptability	9.8

Chemicals characteristics of stored mandarin viz. acidity, ascorbic acid, total sugars and T.S.S. were determined and data presented in [Table-3 and 4].

Titrate acidity

An examination of [Table-3] reveals that the pre-treatments significantly affected the acidity of mandarin. The minimum acidity (0.598%) was recorded in treatment T₀ and the maximum (0.613%) was recorded in T₂ after twenty days of storage. The maximum acidity (0.614%) was recorded in P₂ and minimum (0.596%) in P₀ after twenty days of storage. Acidity of fruits decreased with the advancement of storage period. When all the individual treatments were combined, their effect was found to be significant. Combined application of 2, 4-D 100 ppm with perforated polythene (T₂P₂) resulted in maximum acidity as compared to maximum in control (T₀P₀) at the end of storage period i.e. 20th day. It might be due to low degree of oxidation of organic acids in perforated polythene bags. The present findings are supported by [6-11].

Ascorbic acid

Among various chemical parameter, ascorbic acid or vitamin 'C' is very important qualitative parameter of mandarin fruits. The minimum ascorbic acid (12.45 mg/100 ml juice) was recorded in treatment T₀ and the maximum (13.13 mg/100 ml juice) was obtained in T₂ after twenty days of storage. The maximum ascorbic acid (13.17 mg/100 ml juice) was in P₁ and minimum (11.87 mg/100 ml juice) in P₀ after twenty days of storage. The maximum (14.17 mg/100 ml juice) ascorbic acid was obtained in T₂P₁ and minimum (11.87 mg/100 ml juice) in T₀P₀ after twenty days of storage. In the present study, ascorbic acid content in mandarin fruit was significantly affected by different chemicals, packaging materials and their combinations throughout the storage period up to 20th day. The ascorbic acid content of fruits decreased with advancement of storage period. However, various chemical and packaging materials used in the present study helped in reducing the loss of vitamin 'C' during storage fruits. The decrease in ascorbic acid during storage is probably due to the process of oxidation of ascorbic acid to dehydro-ascorbic acid by enzyme ascorbic-nase. The reduction in the loss of vitamin 'C' content of mandarin fruits due to various chemical and packaging materials treatment as obtained in present study may be due to reduction in the rate of evapo-transpiration, which normally results in volatile dissipation of ascorbic acid during storage. The control (T₀P₀) showed lower retention of ascorbic acid while the higher retention of ascorbic acid content of fruits was observed in 2,4-D 100 ppm + corrugated boxes (T₂P₁). The present findings are supported by [9-11].

Table-3 Effect of pre-treatments, packaging materials and their combination on acidity (%) and ascorbic acid content (mg/100 ml of juice) of mandarin

Treatments	Acidity (%)				Ascorbic acid content (mg/100 ml)			
	5 DAS	10 DAS	15 DAS	20 DAS	5 DAS	10 DAS	15 DAS	20 DAS
T ₀	0.838	0.748	0.668	0.598	15.68	14.38	13.28	12.45
T ₁	0.849	0.759	0.679	0.609	15.91	14.62	13.51	12.67
T ₂	0.853	0.764	0.686	0.613	16.35	15.03	13.95	13.13
T ₃	0.847	0.757	0.677	0.608	15.73	14.43	13.33	12.49
T ₄	0.846	0.756	0.676	0.606	15.72	14.43	13.32	12.49
S.Em±	0.002	0.002	0.002	0.0019	0.04	0.037	0.034	0.027
CD at 5%	0.0055	0.00553	0.007	0.0057	0.12	0.106	0.098	0.078
P ₀	0.836	0.746	0.666	0.596	15.10	13.80	12.70	11.87
P ₁	0.849	0.759	0.679	0.609	16.40	15.10	14.00	13.17
P ₂	0.853	0.764	0.686	0.614	15.70	14.40	13.30	12.50
P ₃	0.849	0.759	0.679	0.609	15.50	14.20	13.10	12.27
S.Em±	0.0017	0.00173	0.0021	0.0018	15.63	14.33	13.23	12.40
CD at 5%	0.0049	0.005	0.0059	0.0051	16.40	15.13	14.00	13.13
T ₀ P ₀	0.821	0.731	0.651	0.581	15.10	13.80	12.70	11.87
T ₀ P ₁	0.845	0.755	0.675	0.605	16.40	15.10	14.00	13.17
T ₀ P ₂	0.847	0.757	0.677	0.607	15.70	14.40	13.30	12.50
T ₀ P ₃	0.840	0.750	0.670	0.600	15.50	14.20	13.10	12.27
T ₁ P ₀	0.839	0.749	0.669	0.599	15.63	14.33	13.23	12.40
T ₁ P ₁	0.854	0.764	0.684	0.614	16.40	15.13	14.00	13.13
T ₁ P ₂	0.853	0.763	0.683	0.613	15.83	14.53	13.43	12.60
T ₁ P ₃	0.849	0.759	0.679	0.609	15.77	14.47	13.37	12.53
T ₂ P ₀	0.838	0.748	0.668	0.598	15.50	14.20	13.10	12.27
T ₂ P ₁	0.856	0.766	0.686	0.616	17.40	16.10	15.00	14.17
T ₂ P ₂	0.864	0.776	0.706	0.624	16.30	14.97	13.90	13.10
T ₂ P ₃	0.855	0.765	0.685	0.615	16.20	14.83	13.80	12.97
T ₃ P ₀	0.837	0.747	0.667	0.597	15.63	14.33	13.23	12.40
T ₃ P ₁	0.845	0.755	0.675	0.605	16.13	14.83	13.73	12.90
T ₃ P ₂	0.856	0.767	0.687	0.621	15.43	14.13	13.03	12.20
T ₃ P ₃	0.849	0.759	0.679	0.609	15.30	14.00	12.90	12.07
T ₄ P ₀	0.844	0.754	0.674	0.604	15.70	14.40	13.30	12.43
T ₄ P ₁	0.845	0.755	0.675	0.605	16.37	15.10	13.97	13.13
T ₄ P ₂	0.846	0.756	0.676	0.606	15.43	14.13	13.03	12.20
T ₄ P ₃	0.850	0.760	0.680	0.610	15.37	14.07	12.97	12.20
S.Em±	0.0038	0.0039	0.0047	0.0040	0.075	0.074	0.068	0.054
CD at 5%	0.0109	0.0111	0.0134	0.0113	0.215	0.211	0.195	0.155

S.Em±: Standard error of mean, CD: Critical difference, DAS: Days After Storage

Table-4 Effect of pre-treatments, packaging materials and their combination on TSS (° Brix) and Total sugars (%) of mandarin

Treatments	TSS (° Brix)				Total sugars (%)			
	5 DAS	10 DAS	15 DAS	20 DAS	5 DAS	10 DAS	15 DAS	20 DAS
T ₀	9.31	10.67	11.96	12.60	8.05	8.85	9.46	9.95
T ₁	8.70	10.06	11.35	12.00	7.90	8.70	9.30	9.80
T ₂	8.25	9.61	10.88	11.55	7.83	8.62	9.22	9.72
T ₃	8.58	9.94	11.22	11.87	7.93	8.73	9.33	9.83
T ₄	8.59	9.91	11.24	11.88	7.87	8.67	9.27	9.77
S.Em±	0.0265	0.0337	0.0295	0.0267	0.047	0.048	0.047	0.047
CD at 5%	0.0757	0.0963	0.0843	0.0762	0.136	0.136	0.136	0.136
P ₀	9.22	10.55	11.87	12.52	8.00	8.80	9.41	9.90
P ₁	8.67	10.03	11.30	11.97	7.88	8.68	9.28	9.78
P ₂	8.18	9.54	10.83	11.47	7.83	8.62	9.22	9.72
P ₃	8.67	10.03	11.31	11.95	7.96	8.76	9.36	9.86
S.Em±	0.0237	0.0301	0.0264	0.0238	0.0425	0.0425	0.0424	0.1212
CD at 5%	0.0677	0.0861	0.0754	0.0681	0.1214	0.1215	0.1212	0.1214
T ₀ P ₀	9.82	11.18	12.47	13.12	8.18	8.98	9.61	10.08
T ₀ P ₁	9.11	10.47	11.76	12.41	8.07	8.87	9.47	9.97
T ₀ P ₂	8.99	10.35	11.64	12.29	7.92	8.72	9.32	9.82
T ₀ P ₃	9.30	10.66	11.95	12.56	8.05	8.85	9.45	9.95
T ₁ P ₀	9.03	10.39	11.68	12.33	7.99	8.79	9.39	9.89
T ₁ P ₁	9.02	10.38	11.67	12.32	7.77	8.57	9.17	9.67
T ₁ P ₂	8.04	9.40	10.69	11.34	7.86	8.66	9.26	9.76
T ₁ P ₃	8.69	10.05	11.34	11.99	7.97	8.77	9.37	9.87
T ₂ P ₀	9.03	10.39	11.68	12.33	7.73	8.53	9.13	9.63
T ₂ P ₁	8.15	9.51	10.71	11.45	7.85	8.65	9.25	9.75
T ₂ P ₂	7.65	9.01	10.30	10.95	7.69	8.46	9.06	9.56
T ₂ P ₃	8.17	9.53	10.82	11.47	8.04	8.84	9.44	9.94
T ₃ P ₀	9.10	10.46	11.75	12.40	8.06	8.86	9.46	9.95
T ₃ P ₁	8.53	9.89	11.18	11.83	7.89	8.69	9.29	9.79
T ₃ P ₂	8.11	9.47	10.76	11.38	7.84	8.64	9.24	9.74
T ₃ P ₃	8.58	9.94	11.20	11.88	7.95	8.75	9.35	9.85
T ₄ P ₀	9.10	10.30	11.75	12.40	8.05	8.85	9.45	9.95
T ₄ P ₁	8.53	9.89	11.18	11.83	7.83	8.63	9.23	9.73
T ₄ P ₂	8.11	9.47	10.76	11.41	7.84	8.64	9.24	9.74
T ₄ P ₃	8.60	9.96	11.25	11.86	7.77	8.57	9.17	9.67
S.Em±	0.0529	0.0674	0.0590	0.0533	0.0949	0.0950	0.0948	0.0950
CD at 5%	0.1513	0.1926	0.1686	0.1523	0.2713	0.2716	0.2711	0.2714

S.Em±: Standard error of mean, CD: Critical difference, DAS: Days After Storage

Table-5 Effect of pre-treatments, packaging materials and their combination on colour and overall acceptability scores of mandarin

Treatments	Colour				Overall acceptability			
	5 DAS	10 DAS	15 DAS	20 DAS	5 DAS	10 DAS	15 DAS	20 DAS
T ₀	7.99	6.95	4.45	3.02	8.39	7.46	5.05	3.73
T ₁	8.30	7.24	4.74	3.33	8.70	7.75	5.34	4.02
T ₂	8.70	7.65	5.14	3.71	9.10	8.17	5.73	4.40
T ₃	8.12	7.08	4.57	3.15	8.52	7.58	5.17	3.85
T ₄	8.09	7.06	4.55	3.13	8.49	7.56	5.15	3.83
S.E.m±	0.0344	0.0333	0.0323	0.0326	0.0344	0.0310	0.0318	0.0315
CD at 5%	0.0982	0.0953	0.0923	0.0931	0.0982	0.0885	0.0910	0.0901
P ₀	7.96	6.91	4.41	2.99	8.36	7.42	5.01	3.69
P ₁	8.48	7.45	4.93	3.50	8.88	7.95	5.52	4.21
P ₂	8.24	7.19	4.68	3.27	8.64	7.70	5.28	3.96
P ₃	8.27	7.23	4.73	3.30	8.67	7.74	5.33	4.00
S.E.m±	0.031	0.03	0.029	0.0291	0.03	0.0277	0.0285	0.0282
CD at 5%	0.088	0.085	0.0825	0.0833	0.0878	0.0791	0.0814	0.081
T ₀ P ₀	7.82	6.80	4.29	2.87	8.22	7.30	4.89	3.57
T ₀ P ₁	8.22	7.20	4.69	3.27	8.62	7.70	5.29	3.97
T ₀ P ₂	7.95	6.90	4.39	2.97	8.35	7.40	4.99	3.67
T ₀ P ₃	7.95	6.90	4.42	2.97	8.35	7.43	5.02	3.70
T ₁ P ₀	7.96	6.90	4.42	3.00	8.36	7.43	5.02	3.70
T ₁ P ₁	8.32	7.27	4.76	3.34	8.72	7.77	5.36	4.04
T ₁ P ₂	8.39	7.37	4.86	3.46	8.79	7.87	5.46	4.14
T ₁ P ₃	8.52	7.43	4.92	3.50	8.92	7.93	5.52	4.20
T ₂ P ₀	8.05	6.97	4.46	3.04	8.45	7.47	5.06	3.74
T ₂ P ₁	9.26	8.23	5.69	4.24	9.66	8.73	6.26	4.97
T ₂ P ₂	8.72	7.67	5.16	3.74	9.12	8.20	5.76	4.40
T ₂ P ₃	8.77	7.73	5.26	3.84	9.17	8.27	5.86	4.49
T ₃ P ₀	8.01	6.97	4.46	3.04	8.41	7.47	5.06	3.74
T ₃ P ₁	8.39	7.37	4.86	3.44	8.79	7.87	5.46	4.14
T ₃ P ₂	8.02	6.97	4.46	3.04	8.42	7.47	5.06	3.74
T ₃ P ₃	8.05	7.00	4.49	3.07	8.45	7.50	5.09	3.77
T ₄ P ₀	7.98	6.93	4.42	3.00	8.38	7.43	5.02	3.70
T ₄ P ₁	8.22	7.17	4.66	3.24	8.62	7.67	5.26	3.94
T ₄ P ₂	8.11	7.07	4.56	3.14	8.51	7.57	5.16	3.84
T ₄ P ₃	8.07	7.07	4.56	3.14	8.47	7.57	5.16	3.84
S.E.m±	0.0687	0.0667	0.0646	0.0652	0.0687	0.0619	0.0637	0.0630
CD at 5%	0.1964	0.1906	0.1845	0.1863	0.1964	0.1770	0.1820	0.1801

Total soluble solids (TSS)

Minimum TSS (11.55° Brix) was recorded in treatment T₂ and the maximum (12.60° Brix) was recorded in T₀ after twenty days of storage. TSS (° Brix) of stored mandarin was significantly affected by packaging methods. The maximum TSS (12.52° Brix) was recorded in P₀ and minimum (11.47° Brix) in P₂ after twenty days of storage. The maximum (13.12° Brix) TSS (° Brix) was obtained in T₀P₀ and minimum (10.95° Brix) in T₂P₂ after twenty days of storage. It is clear from the data that TSS content of fruits was increased with advancement of storage period. The minimum increase in TSS of fruits during storage period is desirable for preservation of good fruit quality. The minimum TSS content was observed in 2,4-D 50 ppm + perforated polythene (T₂P₂) on 20th day of storage, however it was maximum in control (T₀P₀). [12] reported that the rate of increase in TSS was slower in perforated polythene as compared to no packaging. Increase in TSS with advancement of storage may be accounted to the moisture loss, hydrolysis of polysaccharides and concentration of juice as a result of dehydration. The present findings are supported by [13].

Total sugars

A minimum total sugar content of 9.56% was obtained in T₂P₂ and maximum (10.80%) in T₀P₀ after twenty days of storage. It is comprehensible from the data that total sugar of fruits was increased with advancement of storage period. Combined effect of chemical and packaging materials was found to be non-significant during the storage period. The minimum content of total sugars was observed in 2,4-D 100 ppm + perforated polythene (T₂P₂) on 20th day of storage, however it was maximum in control (T₀P₀). The effectiveness of application of different chemicals, packaging materials and their combination might be due to fact that some acids being converted into sugars during respiration conversion of starch (polysaccharides) into sugars (monosaccharide). The present findings are supported by [9-11].

Organoleptic characteristics

Sensory parameters such as fruit colour, texture, flavor and overall acceptability scores were maximum for the fruits treated with fruit treated with 100 ppm 2,4-D + corrugated boxes (T₂P₁). Maximum score for colour (4.24) was obtained in T₂P₂ and minimum (2.87) in T₀P₀ after twenty days of storage. Maximum colour retention by T₂P₁ might be due to inhibition of chlorophyll conversion into carotenoids, lycopene and β-carotene [14, 15]. Marketability of the produce depends on overall acceptability and it was highest for T₂P₁. Maximum value for overall acceptability was obtained in T₂P₁ (4.97) and minimum in T₀P₀ (3.57) after twenty days of storage. This might be due to the fact that packaging prevents the direct evapo-transpiration and lowered the physiological loss in weight [11, 16, 17].

Conclusion

The combined application of 2, 4-D 100 ppm with perforated polythene (P₂T₂) proved to be best post harvest treatment for storage of mandarin from the point of fruit quality in terms of total soluble solids (TSS), acidity and total sugars content during storage on 20th day of storage. Therefore, it may be recommended that before transportation, growers and retailers shall pack the mandarin fruits in perforated polythene with 2.5 mm holes after the chemical treatment with 2, 4-D 100 ppm.

Conflict of Interest: None declared**References**

- [1] Sonkar R.K. and Ladaniya M.S. (1995) *Indian Food Packer*, 49 (6), 11-16.
- [2] Ladaniya M.S. and Sonkar R.K. (1996) *Indian J. Agric. Sci.*, 66, 109-113.
- [3] Ranganna S. (1978) *Manual of analysis of fruit and vegetable products*. Tata McGraw Hill Publisher, New Delhi.
- [4] A.O.A.C. (1960) *Official Methods of Analysis*, Association of the Official

Analytical Chemists, Washington, D.C. 8th Edn.

- [5] Fisher R.A. (1950) *Statistical Inference and Analysis*, Oxford University Press, London.
- [6] Bhullar J.S., Farmahan H.L. and Agnihotri R.P. (1981) *Prog. Hort.*, 13 (3-4), 115-119.
- [7] Tatlı H. and Ozguven A.I. (1999) *Turkish Journal of Agriculture & Forestry*, 23 (supplement 5), 1033-1042.
- [8] Jadhao S.D., Borkar P.A., Bakane P.H., Shinde K.J. and Murumkar R.P. (2008) *Journal of Soils and Crops*, 18 (2), 422-427.
- [9] Ahmad M., Khalid Z.M. and Farooqi W.A. (1979) *Proc. Fla. State Hort. Soc.*, 92, 237-240.
- [10] Ahmad M. and Khan I. (1987) *Plant Foods Human Nutr.*, 37, 47-57.
- [11] Ladaniya, M.S. (2001) *Indian J. Agric. Sci.*, 71, 234-239.
- [12] Deka B.C., Sharma S. and Borah S.C. (2006) *Indian J. Hort.*, 63(3), 251-255.
- [13] Jadhao S.D., Borkar P.A., Borkar S.L., Bakane P.H. and Murumkar R.P. (2008) *Asian Journal of Bio-Science*, 3(2), 247-250.
- [14] Siddiqui S., Sharma R.K. and Gupta O.P. (1997) *Acta Horticulturae*, 735, 535-545.
- [15] Mandhyan B.L. (1999) *Journal of Food Science and Technology*, 36(1), 46-48.
- [16] Sonkar R.K. and Ladaniya M.S. (1999) *Journal of Food Science and Technology*, 36(3), 273-276.
- [17] Choudhary M.R. and Dhaka R.S. (2005) *Haryana Journal of Horticultural Sciences*, 34(1/2), 39-41.